

Non-commutative compactifications and elliptic curves

Yan Soibelman, Vadim Vologodsky

February 1, 2008

To Yuri Ivanovich Manin on his 65th birthday

Abstract. We introduce non-commutative degenerations of elliptic curves over local fields. Corresponding objects are close relatives of non-commutative tori of Connes and Rieffel.

1 Introduction

Let X be an analytic space over a local field ($\mathbf{C}$, $\mathbf{R}$ or $\mathbf{Q}_p$), and G be a group acting freely on X . If the action is not discontinuous the quotient X/G is not an analytic space. Nevertheless one can define the “category of coherent sheaves on X/G ” (notation $Coh(X/G)$) as the category of G -equivariant coherent sheaves on X . A typical example studied in this paper is $X = \mathbf{C}^*$ and $G = q^{\mathbf{Z}}$ acting by dilations $z \mapsto q^n z$. If $|q| \neq 1$ the quotient $\mathbf{C}^*/q^{\mathbf{Z}}$ is an elliptic curve. If $|q| = 1$ the quotient is not even a Hausdorff space. In either case the category $Coh(X/G)$ is equivalent to the category of certain modules over the cross-product of the algebra $\mathcal{O}(\mathbf{C}^*)$ of analytic functions and the group algebra of $q^{\mathbf{Z}}$ (see Section 3). According to the philosophy of non-commutative compactifications proposed in [So1] the compactification of the moduli space of elliptic curves should also contain objects corresponding to the case $|q| = 1$. They can be viewed as non-commutative degenerations of ordinary elliptic curves and called non-commutative (or quantum) elliptic curves. They also can be called analytic quantum tori because in

non-commutative geometry C^∞ -versions of these objects are called quantum or non-commutative tori. The latter were introduced by Alain Connes and studied by Connes and Marc Rieffel in the early 80's (see for ex. [Co]).

Elliptic curves (commutative or not) can be viewed as q -difference analogs of analytic vector bundles with flat connections. In this sense they are a special case of holonomic q -D-modules (cf. [Sab]). There is a vast literature on q -difference equations. It is mainly devoted to the case $|q| \neq 1$.

From the point of view of non-commutative geometry our non-commutative elliptic curves are “bad” quotients. On the other hand, a complex elliptic curve is a C^∞ -manifold (2-dimensional torus) equipped with a complex structure. Hence one has another degeneration: complex structure degenerates into a foliation. The corresponding categories of vector bundles which carry a flat connection were studied in [So1]. It would be interesting to make more precise the relationship between our non-commutative elliptic curves and quantum tori from [So1]. We remark that the picture of [So1] does not have a p -adic analog, while non-commutative elliptic curves exist over non-archimedean local fields as well.

About the contents of the paper. In Section 2 we prove a theorem characterizing algebraically the category of coherent sheaves on certain quotients of Stein varieties. In Section 3 we define degenerate elliptic curves (there $|q| = 1$). This includes both analytic and formal versions. In particular, in Section 3.2 we revisit the results of [BG] concerning the formal case. We compute their Picard groups and obtain some results about irreducible objects in the category of coherent sheaves. We also discuss the analog of the uniformization of an elliptic curve by a complex line. Using this picture we define the moduli space of non-commutative elliptic curves and prove the analog of Rieffel's theorem about $SL_2(\mathbf{Z})$ -action on the moduli space of quantum tori. Our approach should work in the case of abelian varieties as well. In this case we expect that the analog of the Rieffel-Schwarz theorem holds (see [RS]). The reader will notice that in Section 3 we do not distinguish between ordinary and degenerate elliptic curves calling all of them non-commutative. In the last Section 4 we sketch the case of non-archimedean quantum tori. There is an evidence that using the ideas of [KoSo1] one can apply them to quantization of hyper-Kähler manifolds.

We do not discuss in the paper the general philosophy of non-commutative compactifications as well as various connections and ramifications of this concept. Interested reader can consult with [So1]. There are many speculative relations of non-commutative compactifications with some structures in

number theory, dynamical systems and physics (for ex. with [BDar], [CDS], [De1-2], [Mo1-2]). We hope to address some of them in the future.

Acknowledgements. We thank to Ofer Gabber, Victor Ginzburg, Maxim Kontsevich, Yuri Manin, for interesting discussions. Both authors are thankful to IHES for hospitality and excellent research atmosphere. Y.S. gratefully acknowledges financial support him as a Fellow of the Clay Mathematics Institute.

2 Coherent sheaves on elliptic curves

Let K be either a field of complex numbers or non-archimedean field, $q \in K^*$, $|q| < 1$. Recall that the smooth analytic space $E_q(K) = G_m(K)/q^{\mathbb{Z}} = K^*/q^{\mathbb{Z}}$ is an elliptic curve (in particular it is an algebraic variety). We denote by $Coh(E_q)$ the category of algebraic coherent sheaves on E_q . Since $G_m(K) = K^*$ is an analytic space we have a sheaf of analytic functions $\mathcal{O}_{K^*}$, and the category of analytic coherent sheaves $Coh(K^*)$. The natural projection $\pi : K^* \rightarrow E_q$ gives rise to a faithful exact functor (pull-back) $\pi^* : Coh(E_q) \rightarrow Coh(K^*)$. The image $\mathcal{C}_q = \pi^*(Coh(E_q))$ is the category of $q^{\mathbb{Z}}$ -equivariant analytic coherent sheaves on K^* .

The following proposition is well-known (see [Ki] for non-archimedean version).

Proposition 1 *If $F \in Coh(K^*)$ then $H^i(K^*, F) = 0$ for $i > 0$, and each $F_x, x \in K^*$ is generated by $H^0(K^*, F)$.*

In particular the functor $\phi : V \mapsto H^0(K^*, \pi^*(V))$ is faithful and exact from the category $Coh(E_q)$ to the category of modules over the algebra A_q^{an} generated by the analytic functions $f \in \mathcal{O}(K^*)$ and the invertible generator ξ such that $\xi f(z) = f(qz)\xi$ for all $z \in K^*$. Indeed the space of global sections of an equivariant sheaf on K^* gives rise to an A_q^{an} -module. Notice that A_q^{an} contains two commutative subalgebras: the algebra $\mathcal{O}(K^*)$ of analytic functions and the algebra $K[q^{\mathbb{Z}}] := K[\xi, \xi^{-1}]$.

The aim of this section is to prove the following result.

Theorem 1 *The functor ϕ provides an equivalence of $Coh(E_q)$ with the category $\mathcal{B}_q$ of such A_q^{an} -modules M that M is finitely presentable over $\mathcal{O}(K^*)$, i.e. there exists $m, n > 0$ and an exact sequence of $\mathcal{O}(K^*)$ -modules*

$$\mathcal{O}^n(K^*) \rightarrow \mathcal{O}^m(K^*) \rightarrow M \rightarrow 0.$$

This theorem can be generalized to abelian varieties, and furthermore to certain class of Stein spaces. We would like to state and prove such a generalization.

Let X be a Stein space over K , and G be a group freely acting on X . We assume that X/G is a projective algebraic variety and the pull-back $\pi^*(\mathcal{O}_{X/G}(1))$ is a trivial analytic line bundle on X (here $\pi : X \rightarrow X/G$ is the natural projection). The functor π^* gives rise to an equivalence of the category $Coh(X/G)$ of algebraic coherent sheaves on X/G and the category $Coh_G(X)$ of G -equivariant analytic coherent sheaves on X . The functor of global sections $\Gamma_X : F \rightarrow \Gamma(X, F)$ is an exact functor from $Coh_G(X)$ to the category of modules over the algebra $\mathcal{O}(X, G) := \mathcal{O}(X) \rtimes G$ (cross-product of G and the algebra of analytic functions $\mathcal{O}(X)$). We would like to describe the image of $Coh(X/G)$ under the functor $\phi = \Gamma_X \circ \pi^*$. In order to do that we introduce the category $\mathcal{B}_G(X)$ of $\mathcal{O}(X, G)$ -modules M which are finitely presentable over $\mathcal{O}(X)$. This means that there exist $m, n > 0$ and an exact sequence of $\mathcal{O}(X)$ -modules

$$\mathcal{O}^n(X) \rightarrow \mathcal{O}^m(X) \rightarrow M \rightarrow 0.$$

Theorem 2 *The functor ϕ gives rise to an equivalence of categories*

$$Coh(X/G) \simeq \mathcal{B}_G(X)$$

Clearly the previous theorem follows from this one for $X = K^*$, $G = \mathbf{Z}$ acting by $n(x) = q^n x$, $n \in \mathbf{Z}$, $x \in K^*$.

Proof. The proof will consists of several steps.

Step 1. Let us check that if $F \in Ob(Coh(X/G))$ then $\phi(F) \in Ob(\mathcal{B}_G(X))$. It follows from the Serre's theorem that there exist vector bundles W_1, W_2 and a morphism $W_1 \rightarrow W_2 \rightarrow F \rightarrow 0$. Moreover, we can choose $W_k = \bigoplus_i \mathcal{O}(m_i)$, $k = 1, 2$.

Pulling back W_i , $i = 1, 2$ to X we get a G -equivariant analytic bundles $\pi^*(W_i)$, $i = 1, 2$ and G -equivariant analytic coherent sheaf $\pi^*(F)$. The functor ϕ is exact as the composition of two exact functors (Γ_X is exact because X is Stein). Thus we have an exact sequence of $\mathcal{O}(X)$ -modules

$$\phi(W_1) \rightarrow \phi(W_2) \rightarrow \phi(F) \rightarrow 0.$$

By our assumption the pull-backs of the sheaves $\mathcal{O}_{X/G}(l)$, $l \in \mathbf{Z}$ are trivial analytic vector bundles on X . Hence $\phi(W_i)$, $i = 1, 2$ are free $\mathcal{O}(X)$ -modules of finite ranks. It follows that $\phi(F)$ is an object of the category $\mathcal{B}_G(X)$.

Step 2. Observe that if $F \neq 0$ then $\phi(F) \neq 0$. Indeed, $\pi^*(F) \neq 0$ because π^* is a faithful functor. Since X is Stein the sheaf $\pi^*(F)$ is generated by its global sections which is $\phi(F)$. Hence if $\phi(F) = 0$ then $F = 0$. Similarly one proves that if $f : V \rightarrow W$ is a non-trivial morphism of coherent sheaves on X/G then $\phi(f) \neq 0$. Indeed, consider the subsheaf $Im(f)$ of W . Then $\pi^*(Im(f))$ is generated by $\Gamma(X, \pi^*(Im(f))) \simeq Im(\phi(f))$.

Thus we have proved that $\phi : Coh(X/G) \rightarrow \mathcal{B}_G(X)$ is an exact faithful functor.

Step 3. We would like to prove that any object of $\mathcal{B}_G(X)$ is isomorphic to some $\phi(F)$, $F \in Coh(X/G)$.

Let M be an object of the category $\mathcal{B}_G(X)$. Then for some $m, n > 0$ we have an exact sequence of $\mathcal{O}(X)$ -modules

$$\mathcal{O}^n(X) \rightarrow \mathcal{O}^m(X) \rightarrow M \rightarrow 0.$$

We consider the localization functor $M \mapsto \widetilde{M} := \mathcal{O}_X \otimes M$, where $\mathcal{O}_X$ is the sheaf of analytic functions on X . The localization functor is not faithful on finitely generated modules. It is easy to construct an example of a module $V \neq 0$ such that $\widetilde{V} = 0$. Nevertheless we have the following result.

Lemma 1 *The localization functor gives rise to an equivalence of the category of finitely representable $\mathcal{O}(X)$ -modules and the full subcategory of coherent sheaves whose objects are sheaves isomorphic to $coker(\mathcal{O}_X^n \xrightarrow{f} \mathcal{O}_X^m)$ for some n, m and f .*

The inverse functor Γ_X associates to a sheaf F the $\mathcal{O}(X)$ -module of its global sections.

The lemma follows from the exactness of the localization functor and the functor Γ_X and from fact that $\mathcal{O}^n(X) = \mathcal{O}_X^n$, $\Gamma_X(\mathcal{O}_X^n) = \mathcal{O}^n(X)$.

Returning to the proof of the theorem we see that

$$\Gamma_X(\widetilde{F}) = F$$

for any $F \in Coh(X/G)$. Summarizing, we have constructed an exact fully faithful functor ϕ between the categories in question. This completes the proof. ■

We do not know whether Theorem 2 holds without the assumption of triviality of the pull-back of $\mathcal{O}_{X/G}(1)$.

3 Elliptic curves for $|q| = 1$

3.1 Category $\mathcal{B}_q$

Let K be either the field $\mathbf{C}$ or a non-archimedean local field. Let X be Stein space over K , $\mathcal{O}(X)$ algebra of analytic functions on X , G a group acting freely on X .

Definition 1 *The category of coherent sheaves on the non-commutative space X/G is defined as a category of modules over the cross-product algebra $\mathcal{O}(X) \rtimes G$ which are finitely presentable over $\mathcal{O}(X)$.*

From the point of view of this definition one should not distinguish between “good” and “bad” quotients. They all are non-commutative spaces.

Let us assume that $K = \mathbf{C}$. We will use the above definition in two cases: $X = \mathbf{C}^*, G = q^{\mathbf{Z}}$ and $X = \mathbf{C}, G = \mathbf{Z} \oplus \mathbf{Z}$ with the natural actions of G (by dilations in the first case and by shifts by the lattice in the second one). In this section we will study the case $|q| = 1$ or when the action of the lattice on $\mathbf{C}$ is not discontinuous.

By the very definition the category $\mathcal{B}_1$ is the category of coherent sheaves on $\mathbf{C}^*$, whose space of global sections is finitely presentable over $\mathcal{O}(\mathbf{C}^*)$, endowed with an automorphism.

It is likely that the category $\mathcal{B}_q$ is also equivalent to the latter, provided that q is a root of unity. We do not prove this fact here¹ and will not use it.

From now on we assume that $|q| = 1$ and q is not a root of 1. We start with the following lemma.

Lemma 2 *Let $V \in \text{Ob}(\mathcal{B}_q)$. Then the corresponding coherent sheaf $\tilde{V}$ is free of finite rank.*

Proof. The sheaf $\tilde{V}$ is isomorphic to $\text{coker}(\mathcal{O}_{\mathbf{C}^*}^n \xrightarrow{f} \mathcal{O}_{\mathbf{C}^*}^m)$ for some n, m and f . Let k be the maximal rank of $f_x : \mathbf{C}^n \rightarrow \mathbf{C}^m$ for $x \in \mathbf{C}^*$ and S be the subset of $\mathbf{C}^*$ where the rank is strictly smaller the k . Since S is contained in the set of zeros of a holomorphic function on $\mathbf{C}^*$ (a minor of the matrix f), it is discrete. On the other hand, S must be $\mathbf{Z}$ -invariant. Hence S is empty i.e. $\tilde{V}$ is a vector bundle. It remains to observe that any holomorphic vector bundle on $\mathbf{C}^*$ is trivial.

¹Reader may wish to look at the beginning of Section 3.2 where a similar result is proven.

Proposition 2 *The category $\mathcal{B}_q$ is abelian for all $q \neq 0$.*

Proof. For any object V of $\mathcal{B}_q$ we have the corresponding coherent sheaf $\widetilde{V}$ on $\mathbf{C}^*$. If $f : V \rightarrow W$ is a morphism in $\mathcal{B}_q$ then $\widetilde{Ker(f)}$ and $\widetilde{Coker(f)}$ are coherent sheaves. According to the previous lemma they are both free sheaves of finite rank. Hence the spaces of sections $Ker(f)$ and $Coker(f)$ are A_q^{an} -modules which are of the type $\mathcal{O}^p(\mathbf{C}^*)$ as $\mathcal{O}(\mathbf{C}^*)$ -modules. Hence they belong to $\mathcal{B}_q$. ■

Let $K(\mathbf{C}^*)_{mer}$ be the field of meromorphic functions on $\mathbf{C}^*$. Then we have a functor $\mu : V \mapsto V \otimes_{\mathcal{O}(\mathbf{C}^*)} K(\mathbf{C}^*)_{mer}$ from the category $\mathcal{B}_q$ to the category of finite-dimensional $K(\mathbf{C}^*)_{mer}$ -vector spaces.

Proposition 3 *The functor μ is fully faithful.*

Proof. Every object of $\mathcal{B}_q$ gives rise to a coherent sheaf on $\mathbf{C}^*$. Hence it suffices to prove the corresponding statement for coherent sheaves. It is easy to see that if $V \neq 0$ then $\mu(V) \neq 0$ (otherwise V is a torsion analytic sheaf which is $q^{\mathbf{Z}}$ equivariant. This is impossible because orbits of the $q^{\mathbf{Z}}$ -action are dense in circles $|z|=\text{const}$). Applying this observation to the kernel of a morphism we see that μ is a monomorphism on morphisms. ■

Notice that the functor ξ is not fully faithful in the case $|q| \neq 1$: it kills all torsion sheaves.

Remark 1 *The above results hold for $K = \mathbf{Q}_p$ as well.*

Proposition 4 *Suppose that q does not satisfy the Liouville property, i.e. there exists $L > 0$ such that $|q^n - 1| = O(n^{-L})$. Then the Picard group (i.e. the of isomorphism classes of objects of $\mathcal{B}_q$ which are rank one modules over $\mathcal{O}(\mathbf{C}^*)$) is isomorphic to an extension of $\mathbf{Z}$ by $\mathbf{C}^*/q^{\mathbf{Z}}$.*

Proof. Let $A = \mathcal{O}(\mathbf{C}^*)$. Then A is a $\mathbf{Z}$ -module: the generator $1 \in \mathbf{Z}$ sends $f(z)$ to $f(qz)$. Clearly the Picard group is isomorphic to $H^1(\mathbf{Z}, A^*)$. The proposition follows easily from the exact sequence

$$0 \rightarrow \mathbf{Z} \rightarrow A \xrightarrow{exp} A^* \xrightarrow{v} \mathbf{Z} \rightarrow 0$$

(here $v(f) = \int d \log f = 0$ where the integral is taken over any small circle around $z = 0$) and from the following lemma

Lemma 3 *Suppose that q does not satisfy the Liouville property. Then $H^1(\mathbf{Z}, A) = \mathbf{C}$*

Proof. Let $g(z) = \sum_{-\infty}^{+\infty} g_n z^n$. If $g_0 = 0$ then the equation $g(z) = a(qz) - a(z)$ has a solution $a(z) = \sum_{-\infty}^{+\infty} a_n z^n$, where $a_0 = 0, a_n = g_n/(q^n - 1), n \neq 0$. If q does not satisfy the Liouville property the series converges. ■

Remark 2 *It follows from the proof above that for any q , which is not a root of 1 there is a canonical surjective homomorphism:*

$$\deg : \text{Pic}(\mathcal{B}_q) \rightarrow \mathbf{Z}$$

The number $\deg(V)$ is called the degree of the object V .

It is an interesting problem to describe simple objects of the category $\mathcal{B}_q, |q| = 1$. In the case $|q| \neq 1$ it is equivalent to finding all irreducible sheaves on the elliptic curve E_q . They correspond to points of E_q (torsion sheaves). A vector bundle is not a simple object in the category of coherent sheaves. The situation changes in the case $|q| = 1$. It follows from Lemma 3 that if $|q| = 1$ and q is not a root of 1, there are no “torsion sheaves” at all. On the other hand, objects of rank 1 over $\mathcal{O}(\mathbf{C}^*)$ are simple.

We are going to construct some simple objects of higher ranks. For general q and s with $s^n = q$ we have the obvious n -sheets coverings $f_n : \mathbf{C}^*/s^{\mathbf{Z}} \rightarrow \mathbf{C}^*/q^{\mathbf{Z}}$ and $g_n : \mathbf{C}^*/q^{\mathbf{Z}} \rightarrow \mathbf{C}^*/s^{\mathbf{Z}}$. The coverings give rise to the functors

$$f_{n*}, g_n^* : \mathcal{B}_s \rightarrow \mathcal{B}_q, g_{n*}, f_n^* : \mathcal{B}_q \rightarrow \mathcal{B}_s$$

Mimicking the usual construction one can define the above functors for *any* $q \in \mathbf{C}^*$. The functor f_{n*} corresponds to the following homomorphism of algebras

$$A_q^{an} \rightarrow A_s^{an}, f(z) \mapsto f(z^n), \xi \rightarrow \xi,$$

while the functor g_n^* corresponds to the homomorphism

$$A_q^{an} \rightarrow A_s^{an}, f(z) \mapsto f(z), \xi \rightarrow \xi^n.$$

The other two functors are defined as the tensor product $A_q^{an} \otimes_{A_s^{an}} V$, where in the case of f_{n*} the A_s^{an} -module structure on A_q^{an} is given by the first above homomorphism of algebras while in the case of g_n^* it is given by the second one. One can prove that pull-back and push-forward functors satisfy the usual properties.

Remind (see the proof of Proposition 4) that there is a canonical morphism from $\mathcal{O}(\mathbf{C}^*)$ to the Picard group of $\mathcal{B}_s$. Given integers k and n with $n > 0$ we denote by $P_{k,1}$ the rank one module corresponding to the function z^k and by $P_{k,n}$ the push-forward $f_{n*}P_{k,1}$. The number k coincides with the degree $\deg(P_{k,n})$.

Lemma 4 *Assume that $|q| = 1$ and q is not a root of unity. Then, for any integer k and $n > 0$ such that $(k; n) = 1$, the module $P_{k,n}$ is simple.*

Proof. Assume the contrary, and choose a proper submodule $W \subset P_{k,n}$. Consider the pull back $f_n^*(W) \subset f_n^*(P_{k,n})$. It is easy to show that $f_n^*(P_{k,n})$ is isomorphic to the direct sum of rank 1 objects of degree k , namely

$$f_n^*(P_{k,n}) = P_{k,1} \otimes (\oplus_{\mu, \mu^n=1} A_s^{an}/(\xi - \mu)),$$

where as usual we denote by $(\xi - \mu)$ the right ideal generated by the expression in the brackets.

Since all objects $P_{k,1} \otimes A_s^{an}/(\xi - \mu)$ are simple and pairwise non-isomorphic it follows that $f_n^*(W)$ is isomorphic to the direct sum of rank 1 objects of the form $P_{k,1} \otimes A_s^{an}/(\xi - \mu)$. In particular, $\deg(\det(f_n^*(W))) = kl$, where $0 < l = rk(W) < n$. On the other hand, $\deg(\det(f_n^*W)) = \deg(f_n^*(\det W)) = n \deg(\det W)$. This contradiction completes the proof. ■

Complete description of simple objects is not known.

Question 1 *How to describe all simple objects of $\mathcal{B}_q$ in the case $|q| = 1$?*

At the end of this subsection we will make few comments about the tensor structure on the category $\mathcal{B}_q$ (for arbitrary ground field K). Let us fix $q \in K^*$. Notice that the category of A_q^{an} -modules is a K -linear symmetric monoidal category. It follows from the interpretation of objects of $\mathcal{B}_q$ as coherent sheaves that the tensor product of finitely presentable over $\mathcal{O}(K^*)$ -modules is again a finitely presentable module. Hence $\mathcal{B}_q$ is a symmetric monoidal category. It can be considered as a difference analog of the category of analytic vector bundles on K^* equipped with a flat connection.

3.2 Formal analog of the category $\mathcal{B}_q$

Let $F := \mathbf{C}((z))$ and $R := \mathbf{C}[[z]]$. We denote by $\mathcal{B}_q^{al}$ the category of finite-dimensional vector spaces over F equipped with a semi-linear automorphism

$\xi : V \rightarrow V$, $\xi f(z)v = f(qz)\xi v$. In other words, this is a category of finite-dimensional representations of the algebra A_q^{al} of non-commutative Laurent polynomials $F[\xi, \xi^{-1}]$.

Lemma 5 *Assume that q is a root of unity. Then the category $\mathcal{B}_q^{al}$ is equivalent to the category of finite-dimensional F -vector spaces endowed with an automorphism.*

Proof. Assume that q is a primitive root of unity of order n (i.e. $q^n = 1$ and $q^k \neq 1$, for $k < n$). It is easy to see that the center of the algebra A_q^{al} is $\mathbf{C}((z^n))[\xi^n, \xi^{-n}] \subset F[\xi, \xi^{-1}]$. Localizing we make A_q^{al} into a sheaf of algebras over the $\text{Spec}(\mathbf{C}((z^n))[\xi^n, \xi^{-n}]) = \mathbb{G}_m$. One can easily check that A_q^{al} is an Azumaya algebra over $\mathbb{G}_m$. An object of $\mathcal{B}_q^{al}$ gives rise to a sheaf of modules over the Azumaya algebra supported on a closed subscheme $S \subset \mathbb{G}_m$ finite over $\text{Spec}(F)$. The Brauer group of F is trivial. Hence the restriction of the Azumaya algebra A_q^{al} to the ind-scheme $\varinjlim S$ (the limit is taken over all finite closed subschemes $S \subset \mathbb{G}_m$) splits, i.e. isomorphic to the matrix algebra. It remains to notice that the category of modules over a split Azumaya algebra is equivalent to the category of $\mathcal{O}$ -modules on the underlying scheme. ■

In what follows we assume that q is not a root of unity.

Following [BG] we call an object V of $\mathcal{B}_q^{al}$ integral if there exists a $\mathbf{C}[\xi, \xi^{-1}]$ -invariant R -lattice $L \subset V$. Integral objects form an abelian subcategory $\mathcal{B}_q^f$ of $\mathcal{B}_q^{al}$. The following theorem was proved in [BG].

Theorem 3 *The category $\mathcal{B}_q^f$ is equivalent to the category of finite-dimensional $\mathbf{C}^*/q^{\mathbf{Z}}$ -graded complex vector spaces endowed with a nilpotent operator which preserves the grading.*

Let us describe a construction of the equivalence functor. It is based on the following easy lemma, proof of which is left to the reader.

Lemma 6 *Let V be an object of $\mathcal{B}_q^{al}$, $L \subset V$ be a $\mathbf{C}[\xi]$ -invariant R -lattice (we do not require L to be ξ^{-1} -invariant), and $\overline{P} \subset L/zL := \overline{L}$ be a $\mathbf{C}[\xi]$ -invariant subspace such that for any eigenvalue α of $\xi : \overline{P} \rightarrow \overline{P}$ the number $q^n \alpha$ is not an eigenvalue of ξ on $\overline{L}$ for all $n \neq 0$. Then the embedding $\overline{P} \hookrightarrow \overline{L}$ has a $\mathbf{C}[\xi]$ -equivariant lifting $P \hookrightarrow L$.*

Given an object V of $\mathcal{B}_q^f$ we can find a $\mathbf{C}[\xi, \xi^{-1}]$ -invariant R -lattice $L \subset V$ such that the eigenvalues of ξ on $\bar{L}$ satisfy the property in the lemma.² It follows that V is isomorphic to $\bar{L} \otimes_{\mathbf{C}} F$ as a $F[\xi]$ -module. The action of ξ on $\bar{L}$ defines a $\mathbf{C}^*$ -grading ($\bar{L}_a := \ker(\xi - a)^n$ for large n). The latter gives rise to a $\mathbf{C}^*/q^{\mathbf{Z}}$ -grading. Finally, we endow $\bar{L}_a$ with a nilpotent operator $\log(a^{-1}\xi)$. It gives rise to the equivalence functor.

Notice that the argument above shows that there is a functor from the category $\mathcal{B}_q^f$ to the category of finitely generated modules over the algebra of non-commutative Laurent polynomials $\mathbf{C}[z, z^{-1}, \xi, \xi^{-1}] \subset A_q^{al}$. The latter sends V to $\tilde{V} = \bar{L} \otimes_{\mathbf{C}} \mathbf{C}[z, z^{-1}]$.

Now we want to obtain a similar description of the whole category $\mathcal{B}_q^{al}$. We claim that for any object V of $\mathcal{B}_q^{al}$, there exists an integer n such that the pull back $f_n^*(V)$ is a direct sum of tensor products of integral objects and one-dimensional ones. More precisely we have the following result.

Theorem 4 *The category $\mathcal{B}_q^{al}$ is equivalent to the category of $\mathbf{Q}$ -graded objects of $\mathcal{B}_q^f$ (i.e. each object of $\mathcal{B}_q^{al}$ is direct sum of $\mathbf{Q}$ copies of objects of $\mathcal{B}_q^f$).*

Proof. Given a rational number $\lambda = \frac{k}{n}$ with $n > 0$, $(n; k) = 1$, $k \neq 0$ unless $n = 1$, we, first, construct a functor

$$G_{k,n} : \mathcal{B}_q^f \rightarrow \mathcal{B}_q^{al}$$

For this purpose we represent the category $\mathcal{B}_q^f$ as a certain “tensor quotient” of the category $\mathcal{B}_s^f$. Namely, consider the full subcategory $\mathcal{E}$ of $\mathcal{B}_s^f$, whose objects are A_s^{al} -modules isomorphic to $A_s^{al}/(\xi - \nu)$, where $\nu \in \mathbf{C}^*$ and $\nu^n = 1$. Notice that $\mathcal{E}$ is a Picard category i.e. a groupoid with a tensor structure such that any object is invertible. Define the “tensor quotient” $\mathcal{B}_s^f/\mathcal{E}$ to be the category whose objects are those of $\mathcal{B}_s^f$ and morphisms between V and W are given by homomorphisms of A_s^{al} -modules $V \otimes M \rightarrow W$

²To construct such a lattice L we can use the following procedure. First choose any $\mathbf{C}[\xi, \xi^{-1}]$ -invariant R -lattice L_0 . Assume, that it does satisfy the eigenvalues property: numbers α and $q^n\alpha$ are eigenvalues, for some $\alpha \neq 0$ and $n > 0$. Let $\bigoplus_{\gamma} \bar{L}_0^{\gamma} = \bar{L}_0 := L_0/zL_0$ be the Jordan decomposition of $\bar{L}_0$ (i.e. $(\xi - \gamma)^N \bar{L}_0^{\gamma} = 0$, for large N). Choose a decomposition $L_0 = \bigoplus_{\gamma} L_0^{\gamma}$ of the R -lattice which descends to the latter one modulo zR . Put $L_1 := \bigoplus_{\gamma \neq q^n\alpha} L_0^{\gamma} \oplus z^{-1}L_0^{q^n\alpha}$. It is easy to see that L_1 is a $\mathbf{C}[\xi, \xi^{-1}]$ -invariant R -lattice. If it still does not satisfy the property in the lemma we apply the above procedure again to L_1 and so on.

where $M = \bigoplus_{\{\nu|\nu^n=1\}} A_s^{al}/(\xi - \nu)$. This is an abelian category. For any object V of $\mathcal{B}_s^f$ and E of $\mathcal{E}$ the object $V \otimes E$ is isomorphic to V . It is easy to see that the functor

$$g_n^* : \mathcal{B}_s^f/\mathcal{E} \rightarrow \mathcal{B}_q^f$$

is an equivalence.

Given an object V of $\mathcal{B}_s^f/\mathcal{E}$ we define $G_{k,n}(V) = f_{n*}(V \otimes P_{k,1})$. (Recall that $P_{k,1}$ stands for $A_s^{al}/(\xi - z^k)$).

We claim that $G_{k,n}$ extends to morphisms thus gives rise to a functor. Indeed,

$$Hom_{\mathcal{B}_q^{al}}(f_{n*}(V \otimes P_{k,1}); f_{n*}(W \otimes P_{k,1})) = Hom_{\mathcal{B}_s^{al}}(f_n^* f_{n*}(V \otimes P_{k,1}); W \otimes P_{k,1})$$

Now the claim is an immediate corollary of the following formula:

$$f_n^* f_{n*}(V \otimes P_{k,1}) \simeq V \otimes P_{k,1} \otimes M \quad (1)$$

We leave its proof to the reader.

Notice, that the latter argument also shows that the functor $G_{k,n}$ is fully faithful.

Next, we are going to show that the functor

$$\bigoplus_{n,m} G_{k,n} : \bigoplus_{k,n} \mathcal{B}_q^f \rightarrow \mathcal{B}_q^{al}$$

is an equivalence of categories.

First, let us prove that the functor is fully faithful. Indeed, it remains to show that

$$Hom_{\mathcal{B}_q^{al}}(G_{k,n}(V), G_{k',n'}(W)) = 0$$

unless $n = n'$ and $k = k'$. Notice, that the group under consideration is a subgroup of

$$Hom(f_{nn'}^* f_{n*}(P_{k,1} \otimes V); f_{nn'}^* f_{n'*}(P_{k',1} \otimes W))$$

The formula (1) easily implies that the latter is trivial.

It remains to show that the image of $\bigoplus_{n,k} G_{k,n}$ is $\mathcal{B}_q^{al}$.

Proposition 5 *For any object V of $\mathcal{B}_q^{al}$ there exists $n > 0$ such that $f_n^*(V)$ is a direct sum of $A_q^{al}/(\xi - az^k)^l$ where $k, l > 0$ are integers and $a \in \mathbb{C}$.*

Proof. We start with the following lemma.

Lemma 7 *All one-sided ideals of A_q^{al} are principal.*

Proof of the Lemma. For an element $X \in A_q^{al}$, $X = a_l \xi^l + \dots + a_m \xi^m$ with $a_l, a_m \neq 0$ we define $s(X) = m - l$. It is easy to see that A_q^{al} is an Euclidean domain relative to the function $s(X)$.

It follows that any finite-dimensional A_q^{al} -module is a direct sum of cyclic ones (i.e. those generated by a single element). Hence, it is enough to prove the proposition for $V = A_q^{al}/(X)$, where $X \in A_q^{al}$. We will do it by induction on the dimension of V .

Without loss of generality we may assume that $X = c\xi^l + a_{l-1}\xi^{l-1} + \dots + a_0$ where a_i are Taylor series in z and $c \in \mathbf{C}^*$. Notice that $L := \text{Im}(R[\xi] \rightarrow V)$ is a ξ -invariant R -lattice in V . If $a_s(0) \neq 0$, for some s , the element ξ induces the operator $\bar{\xi} : L/zL \rightarrow L/zL$ which is not nilpotent. Then we can make use of Lemma 6 to reduce the dimension of V . In general, there exist integers n and k such that $c(z^{k/n}\xi)^l + a_{l-1}(z^{k/n}\xi)^{l-1} + \dots + a_0 = z^{lk/n}(c'\xi^l + b_{l-1}\xi^{l-1} + \dots + b_0)$ with $b_i \in \mathbf{C}[[z]]$ and $b_s(0) \neq 0$, for some s , and we can apply Lemma 6 to $P_{-k,1} \otimes f_n^*(V)$. This completes the proof of the proposition.

Returning to the proof of the theorem we notice that the modules $A_q^{al}/(\xi - az^k)^l$ are in the image of $\oplus_{n,k} G_{k,n}$. The second remark is that the latter image is closed under the push-forward functor and under taking subobjects.

Now, given an object V of $\mathcal{B}_q^{al}$ we choose n as in the proposition and consider the canonical morphism

$$f_{n*}f_n^*(V) \rightarrow V$$

It is easy to see that the morphism is surjective. Combining this with the two observations above we see that V is in the image of $\oplus_{n,k} G_{k,n}$. The theorem is proven. ■

Remark 3 1) *The argument given above is just a variant of Manin's proof of the Classification Theorem for Dieudonné modules (see [M4]).*

2) *One can compare our Theorem 4 with a well-known result from the theory of D -modules which says that, for any vector bundle E with a connection on the punctured formal disk, the pull-back of E on some finite covering of the punctured disk is a direct sum of tensor products of D -modules with regular singularities (at the origin) and one-dimensional ones.*³

³We could not find a reference for this result. We are grateful to Roman Bezrukavnikov who explained it to one of us. He, in turn, refers to the lectures by Spencer Bloch at the University of Chicago.

Corollary 1 *Any finite-dimensional A_q^{al} -module is a direct sum of modules of the type $A_q^{al}/(\xi^n - az^k)^l$ where $n > 0, k, l > 0$ are integers and $a \in \mathbf{C}^*$.*

Corollary 2 *For any two complex numbers q and q' (which are not roots of 1) the categories $\mathcal{B}_q^{al}$ and $\mathcal{B}_{q'}^{al}$ are equivalent.*

We would like to describe the relation between the category $\mathcal{B}_q^{al}$ and its analytic counterpart $\mathcal{B}_q$.

First, we construct a tensor functor

$$\mathcal{B}_q^{al} \rightarrow \mathcal{B}_q \quad (2)$$

In the process of proving Theorem 3 we observed that there was a functor from the category $\mathcal{B}_q^f$ to the category of finitely generated modules over $\mathbf{C}[z, z^{-1}, \xi, \xi^{-1}]$. Composing it with the tensor product with A_q^{an} over the subalgebra $\mathbf{C}[z, z^{-1}, \xi, \xi^{-1}] \subset A_q^{an}$ we obtain the functor $\mathcal{B}_q^f \rightarrow \mathcal{B}_q$. It immediately follows from Theorem 4 (and from the construction of the equivalence given in the proof) that this functor extends to (2).

Assume that $|q| \neq 1$. It is easy to see that the image of (2) lies in the subcategory $Bun(E) \subset \mathcal{B}_q$ of vector bundles on $E := \mathbf{C}^*/q^{\mathbf{Z}}$.

Proposition 6 *Assume that $|q| \neq 1$. Then functor $\mathcal{B}_q^{al} \rightarrow Bun(E)$ induces a bijection on the set of isomorphism classes of objects.*

Proof. It follows from the classification of vector bundles on E obtained by M. Atiyah that any indecomposable vector bundle is of the form $A_q^{an}/(\xi^n - az^k)^l$ where $n > 0, k, l > 0$, $(n, k) = 1$, $a \in \mathbf{C}^*$. Therefore, the functor induces a bijection on the set of isomorphism classes of indecomposable objects. The proposition immediately follows from this fact. ■

It was shown in [BG] that the restriction of (2) to the subcategory $\mathcal{B}_q^f$ gives rise to an equivalence between the latter category and the tensor category of semistable vector bundles on $E := \mathbf{C}^*/q^{\mathbf{Z}}$ of degree zero.

Of course the functor $\mathcal{B}_q^{al} \rightarrow Bun(E)$ is not an equivalence, since $Bun(E)$ is not abelian.

We can reformulate Proposition 6 by saying that there is a bijection between the non-abelian cohomology group $H^1(\mathbf{Z}, GL(n, F))$, $F = \mathbf{C}((z))$ and the set of isomorphism classes of n -dimensional vector bundles on E .

Now, let G be any algebraic group over $\mathbf{C}$. Using the Tannakian formalism and Proposition 6 one can easily construct a map from $H^1(\mathbf{Z}, G(F))$ to the

set of isomorphism classes of principal G -bundles on E . We do not know if it is a bijection.

According to [BG] this is true if restrict ourselves to the subset $Im(H^1(\mathbf{Z}, G(R)) \rightarrow H^1(\mathbf{Z}, G(F)))$, on the one hand, and semistable G -bundles of degree zero, on the other hand.

If $|q| = 1$, we can not say much more about (2), except that in this case $\mathcal{B}_q$ has more objects then its algebraic counterpart $\mathcal{B}_q^{al}$. Probably the Corollary 2 is not true for $\mathcal{B}_q$.

3.3 Uniformization by the complex line

We still assume that $K = \mathbf{C}$. Complex elliptic curves can be also represented as quotients $\mathbf{C}/L$ where L is a lattice in $\mathbf{C}$. If L is not discrete then the quotient can be considered as a non-commutative space. By definition (cf. [M2]) a *quasi-lattice* in $L \subset \mathbf{C}^n$ is a free abelian subgroup of $\mathbf{C}^n$ of rank $2n$. If L is discrete in the standard topology then it is called *lattice*.

Let $L \subset \mathbf{C}$ be a quasi-lattice. We define the algebra C_L as the cross-product of the algebra of analytic functions $\mathcal{O}(\mathbf{C})$ and the group algebra of L . It is generated by analytic functions $f(z)$ and invertible generators $e_\lambda, \lambda \in L$ subject to the relations

$$f_1(z)f_2(z) = f_2(z)f_1(z), e_\lambda e_\mu = e_{\lambda+\mu}, e_\lambda f(z) = f(z + \lambda)e_\lambda.$$

We define the category $\mathcal{D}_L$ as the category of C_L -modules which are finitely presentable over $\mathcal{O}(\mathbf{C})$. Let $\tau \in \mathbf{C} \setminus \mathbf{Q}, q = \exp(2\pi i \tau)$. Let us consider a quasi-lattice $L_\tau = \mathbf{Z} \oplus \mathbf{Z}\tau$. The corresponding non-commutative elliptic curve will be denoted by E_τ .

Proposition 7 *The categories $\mathcal{D}_{L_\tau}$ and $\mathcal{B}_q$ are equivalent.*

Proof. The category $\mathcal{D}_{L_\tau}$ can be identified with the category of L_τ -equivariant analytic coherent sheaves on $\mathbf{C}$. Similarly the category $\mathcal{B}_q$ can be identified with the category of analytic $q^{\mathbf{Z}}$ -equivariant sheaves on $\mathbf{C}^*$. Clearly the category of sheaves are equivalent. ■

3.4 Moduli space of quantum elliptic curves

Let K be a local field, as in Section 2.1, $q \in K^*$.

Recall that the category $\mathcal{B}_q$ defines a *non-commutative (or quantum)* elliptic curve E_q . Of course, if $|q| \neq 1$ it is just the usual elliptic curve over K . The question arises when two quantum elliptic curves should be called equivalent. It is natural to expect (cf. [Ga]) that the equivalences are defined by certain functors between the corresponding categories $\mathcal{B}_q$. We do not know such a definition in general. One of the reasons is lack of the general theory of non-commutative analytic spaces. Existing results in non-commutative algebraic geometry (see for ex. [KR]) are not sufficient for our purposes. However, in the case $K = \mathbf{C}$ we can resolve the difficulty by using the uniformization of elliptic curves by the complex line.

Definition 2 *We say that two non-commutative elliptic curves E_L and $E_{L'}$ corresponding to the categories $\mathcal{D}_L$ and $\mathcal{D}_{L'}$ are equivalent if there is a continuous isomorphism of the algebras C_L and $C_{L'}$ (the topology on the algebras is induced by the topology on the algebra of analytic functions $\mathcal{O}(\mathbf{C})$).*

*Moreover we define **groupoid** of non-commutative elliptic curves putting $\text{Isom}(E_L, E_{L'})$ to be the set of all continuous isomorphisms of the algebras C_L and $C_{L'}$ modulo the group of inner automorphisms of C_L .*

Remark 4 *The reader familiar with non-commutative geometry has noticed that the relationship between the algebras C_L and A_q^{an} is similar to the one between the foliation algebra of a torus and the corresponding quantum torus in [Co]. In particular C_L gives rise to a generator-free description of non-commutative elliptic curves.*

Given a quasi-lattice $L \subset \mathbf{C}$ we define a group G_L to be the subgroup of $\mathbf{C}^*$ consisting of all $\alpha \in \mathbf{C}^*$ such that $L = \alpha L$.

Theorem 5 *a) Non-commutative elliptic curves E_L and $E_{L'}$ are equivalent if and only if L and L' are similar (i.e. there exists $\alpha \in \mathbf{C}^*$ such that $L = \alpha L'$).*

b) $\text{Isom}(E_L, E_L)$ is isomorphic to the semi-direct product $(G_L \triangleright \mathbf{C}/L) \triangleright \text{Pic}(E_L)$ (clearly $\mathbf{C}/L$ is a normal subgroup in G_L , and $\text{Pic}(E_L)$ is a normal subgroup in the semi-direct product of the other two groups).

Proof. The proof will consist of two steps.

i) It is easy to see that the group C_L^* of invertible elements in C_L consists of products $f(z)e_\lambda$, $\lambda \in L$ where $f(z)$ is an invertible analytic function.

Let W_L be a vector space consisting of $f \in C_L$ satisfying the following property: for any invertible $a \in C_L$ there exists a constant c such that

$afa^{-1} = f + c$. Let us prove that such $f = f(z)$ is a linear function of z . Indeed if $f = \sum_j f_j(z)e_{\lambda_j}$, then taking $a = e_\mu, \mu \in L$, we see that the derivatives $f'_j(z)$ are double-periodic analytic function on $\mathbf{C}$ (we do not assume that the periods are linearly independent over $\mathbf{R}$). Hence $f_j(z)$ are linear functions in z .

Clearly $i(W_L) = W_{L'}$. By composing i with an automorphism of C_L we can bring it to the form $i(z) = \alpha z$ and $i(L) \subset L'$, where $\alpha \in \mathbf{C}^*$. Since i is continuous we have $i(f(z)) = f(\alpha z)$ for any $f \in \mathcal{O}(\mathbf{C})$.

Notice that $e_\lambda z e_{-\lambda} = \lambda + z$. Since e_λ is invertible we have $i(e_\lambda) = e_{\lambda'} g(z)$ for some invertible analytic function $g(z)$ and $\lambda' \in L'$. Then $i(e_\lambda z e_{-\lambda}) = \lambda + \alpha z$. On the other hand it is equal to $\alpha \lambda' + \alpha z$. It follows that $\alpha L = L'$.

ii) We let the groups G_L and $\mathbf{C}$ act on C_L preserving the lattice L and sending $z \mapsto \alpha z$, $\alpha \in G_L$ and $z \mapsto z + c$, with $c \in \mathbf{C}$. It is easy to see that the second automorphism is inner provided $c \in L$. It gives rise to an injective homomorphism from the semi-direct product G_L and $\mathbf{C}/L$ to $Isom(E_L, E_L)$. As we have just proven, any automorphism of C_L preserves the subspace of linear in z functions. This implies that the image of the latter homomorphism is a normal subgroup. Hence, it is enough to identify the quotient with $Pic(E_L)$.

An element of the quotient can be represented by an automorphism i of C_L identical on the space of functions of z . We have $i(e_\lambda) = f_\lambda e_\lambda$, where f_λ is an invertible analytic function. It is easy to check that f_λ define a 1-cocycle of L with coefficients in $\mathcal{O}^*$. If i is an inner automorphism, the cocycle f_λ is a coboundary. It gives rise to an isomorphism between the quotient and $H^1(L; \mathcal{O}^*) = Pic(E_L)$. This proves the theorem. ■

Corollary 3 *Two non-commutative elliptic curves E_τ and E_μ are equivalent iff $\tau = g(\mu)$ for some $g \in SL_2(\mathbf{Z})$, where $SL_2(\mathbf{Z})$ acts on $\mu \in \mathbf{C} \setminus \mathbf{Q}$ by fractional transformations.*

It follows from the Corollary that the “moduli space” of non-commutative elliptic curves can be identified with the quotient $\overline{D}/SL_2(\mathbf{Z})$ where $\overline{D} = \{z \in \mathbf{C} \mid |z| \leq 1\}$ and $SL_2(\mathbf{Z})$ acts by fractional transformations. Naively, the moduli space consists of the isomorphism classes of algebras C_L together with the equivalence classes of the categories $\mathcal{B}_q$ where q is a root of 1 (these categories are equivalent to $\mathcal{B}_1$). The action of $SL_2(\mathbf{Z})$ is not discontinuous on the boundary circle ∂D . Hence the moduli space is itself a non-commutative space. It can be described in terms of a cross-product algebra. The reader

should compare Corollary 3 with the similar theorem of Rieffel about quantum tori (see [RS] for the higher-dimensional case).

Remark 5 *It follows from the Corollary that if $\tau \in \mathbf{Q}(\sqrt{d})$, where $d \in \mathbf{Z}$ is square-free then the corresponding non-commutative elliptic curve has more autoequivalences than a generic one. If $d < 0$ it is just an elliptic curve with complex multiplication. It was speculated in [M2] that in the case $d > 0$ non-commutative tori can be used for a description of the maximal abelian extension of $\mathbf{Q}(\sqrt{d})$ (Stark conjectures).*

Since our non-commutative elliptic curve can be defined not only over $\mathbf{C}$, the following questions look very interesting.

Question 2 *Assume that we are working over the local field K as in the Section 2.*

1. *When two categories $\mathcal{B}_q$ and $\mathcal{B}_{q'}$ are equivalent?*
2. *What is the group of auto-equivalences of $\mathcal{B}_q$?*

We don't know the answer even in the case $K = \mathbf{C}$, $|q| = 1$.

For usual elliptic curves (i.e. $|q| \neq 1$) the answer is known thanks to the following result.

Theorem 6 *Assume that $|q|, |q'| \neq 1$. Then*

- a) *The categories $\mathcal{B}_q$ and $\mathcal{B}_{q'}$ are equivalent if and only if the corresponding elliptic curves are isomorphic.*
- b) *The group of auto-equivalences of $\mathcal{B}_q$ is generated by the group of automorphisms of the elliptic curve $E_q = \mathbf{C}^*/q^{\mathbf{Z}}$ and $\text{Pic}(E_q)$. The latter acts on $\mathcal{B}_q$ sending a module V to $V \otimes \mathcal{L}$, where $\mathcal{L} \in \text{Pic}(E_q)$.*

*Proof.*⁴ Notice that irreducible objects of $\mathcal{B}_q$ are skyscraper sheaves supported at points of E_q . Therefore, an equivalence $i : \mathcal{B}_q \rightarrow \mathcal{B}_{q'}$ gives rise to a bijection $\tilde{i}$ between the points of E_q and $E_{q'}$, which is a homeomorphism with respect to the Zariski topology. Further, line bundles $\mathcal{L}$ on E_q can be characterised by the property that $\dim(\text{Hom}(\mathcal{L}, F)) = 1$, for any skyscraper sheaf F . Hence i induces a bijection between line bundles on E_q and $E_{q'}$. Therefore it is enough to prove that if $i(\mathcal{O}_{E_q}) = \mathcal{O}_{E_{q'}}$ then the functor i is induced by an isomorphism of the ringed spaces. Let $S \subset E_q(\mathbf{C})$ be a finite subset. We are

⁴The proof bellow has been explained to the authors by Dima Kaledin.

going to construct a morphism of sheaves restricted to the complements of S

$$i_* : \mathcal{O}(E_q \setminus S) \rightarrow \mathcal{O}(E_{q'} \setminus \tilde{i}(S)) \quad (3)$$

Let $\text{Coh}_S(E_q)$ be the category of coherent sheaves supported on S . The quotient category $\mathcal{B}_q/\text{Coh}_S(E_q)$ is equivalent to the category of coherent sheaves on $E_q \setminus S$. Furthermore $\mathcal{O}(E_q \setminus S)$ can be identified with the ring of endomorphism of the identity functor from $\text{Coh}(E_q \setminus S)$ to itself.

The functor i induces an equivalence between $\text{Coh}_S(E_q)$ and $\text{Coh}_{\tilde{i}(S)}(E_{q'})$ which, in turn, gives rise to an equivalence

$$i : \text{Coh}(E_q \setminus S) \rightarrow \text{Coh}(E_{q'} \setminus \tilde{i}(S))$$

The latter defines the isomorphism (3). ■

Corollary 4 *If L is a lattice then the group $\text{Aut}(\mathcal{D}_L)$ of autoequivalences of the category $\mathcal{D}_L$ is isomorphic to the group $\text{Out}(C_L)$ of outer automorphisms of the algebra C_L .*

Question 3 *Is the Corollary 4 true for a quasi-lattice L ?*

If the answer is positive, then we can say that two non-commutative elliptic curves E_L and $E_{L'}$ are equivalent iff the corresponding categories $\mathcal{D}_L$ and $\mathcal{D}_{L'}$ are equivalent. This observation agrees with the Theorem 6.

3.5 Remark about higher genus curves

If a complex curve X has the genus $g \geq 2$ it admits a universal covering by the upper-half plane H . We have $X \simeq H/\Gamma$, where $\Gamma \subset SL_2(\mathbf{Z})$ is a Fuchsian subgroup. Applying Theorem 2 we see that $\text{Coh}(X) \simeq \mathcal{B}_\Gamma(H)$. Now, if we consider a non-discrete embedding of Γ into $SL_2(\mathbf{Z})$, we define $\text{Coh}(H/\Gamma)$ as $\mathcal{B}_\Gamma(H)$. Since the action of Γ is now ergodic, we conclude that all objects of our category are free $\mathcal{O}(H)$ -modules. In other words, there are no non-free coherent sheaves on degenerate curves. All the questions addressed above to non-commutative elliptic curves can be addressed to non-commutative complex curves of higher genus. Hopefully arising structures can be used for algebraic study of Thurston boundary of Teichmüller space.

4 Non-archimedean quantum tori

This section is independent of the rest of the paper.

Let K be a non-archimedean local field, L is a free abelian group of finite rank d , $\varphi : L \times L \rightarrow K$ is a skew-symmetric bilinear form such that $|\exp(2\pi i\varphi(\lambda, \mu))| = 1$ for any $\lambda, \mu \in L$. We denote by $A(T(L, \varphi))$ the algebra of regular functions on the quantum torus $T(L, \varphi)$ (see [So1]). Thus it is an algebra with generators $e(\lambda), \lambda \in L$, subject to the relation

$$e(\lambda)e(\mu) = \exp(2\pi i\varphi(\lambda, \mu))e(\lambda + \mu).$$

There are various completions of $A(T(L, \varphi))$ which appear in the literature. In this section we will use the following one.

Definition 3 *The space $\mathcal{O}(T(L, \varphi))$ of analytic functions on the non-archimedean quantum torus consists of series $\sum_{\lambda \in L} a(\lambda)e(\lambda), a(\lambda) \in K$ such that*

$|a(\lambda)| \rightarrow 0$ as $|\lambda| \rightarrow \infty$ (here $|(\lambda_1, \dots, \lambda_d)| = \sum_i |\lambda_i|$).

Lemma 8 *Analytic functions $\mathcal{O}(T(L, \varphi))$ form a K -algebra.*

Proof. We have $(\sum_{\lambda} a(\lambda)e(\lambda))(\sum_{\mu} b(\mu)e(\mu)) = \sum_{\nu} c(\nu)e(\nu)$, where $c(\nu) = \sum_{\lambda} a(\lambda)b(\nu - \lambda)\exp(2\pi i\varphi(\lambda, \nu - \lambda))$. We need to prove that $\max_{\lambda, \nu \in L} |a(\lambda)||b(\nu - \lambda)|$ approaches zero as $|\nu| \rightarrow \infty$. Since both functions $a(\lambda)$ and $b(\lambda)$ are bounded on L we may assume that λ belongs to a finite subset of L . Then $|a(\lambda)||b(\nu - \lambda)| \leq \text{const}|b(\nu - \lambda)|$ where $\lambda \in L_0$, and L_0 is a finite set. Therefore $|c(\nu)| \rightarrow 0$ as $|\nu| \rightarrow \infty$. ■

Let us fix an isomorphism $L \simeq \mathbf{Z}^d$ and $r = (r_1, \dots, r_d) \in \mathbf{R}_{>0}^d$. We denote by $\mathcal{O}(T(L, \varphi); r)$ the space of series $f = \sum_{\lambda \in \mathbf{Z}^d} a(\lambda)e(\lambda)$ such that $|a(\lambda)|r^\lambda \rightarrow 0$ as $|\lambda| \rightarrow \infty$. Then $\mathcal{O}(T(L, \varphi); r)$ becomes a non-commutative Banach K -algebra with the norm $\|f\| = \max_{\lambda} |a(\lambda)|r^\lambda$. In the case $\varphi = 0$ an epimorphic image of $\mathcal{O}(T(L, \varphi); r)$ in the category of commutative Banach K -algebras is called an affinoid K -algebra (see [Be]). One can use non-archimedean quantum tori in order to define a deformation of the sheaf of analytic functions on the rigid analytic space obtained from degenerating hyper-Kähler manifolds in the spirit of [KoSo1]. To be more precise, such a degenerating family defines a fibration over an affine manifold Y with the fibers being tori (possibly degenerate on the codimension two subvariety

$Y^{sing} \subset Y$). The sheaf of analytic functions is in fact the sheaf on the non-singular part $Y \setminus Y^{sing}$ only. It is an open problem to define it on Y . One can replace series in commuting variables $x_i, 1 \leq i \leq \dim Y$ by the series in variables x_i such that $x_i x_j = q x_j x_i$. Then one obtains the deformation of the sheaf $\mathcal{O}_Y$. Details will appear in the paper of M. Kontsevich and the first author.

References

- [BG] V. Baranovsky, V. Ginzburg, Conjugacy classes in loop groups and G -bundles on elliptic curves. Int. Math. Res. Notices, 15 (1996), 733-751.
- [Be] V. Berkovich, Spectral theory and analytic geometry over non-Archimedean fields. Mathematical Surveys and Monographs, 33. Amer. Math. Soc., 1990.
- [BDar] M. Bertolini, H. Darmon, The p -adic L -functions of modular elliptic curves, preprint, 2000.
- [Bo] F. Boca, Projections in Rotation Algebras and Theta Functions, math.OA/9803134
- [BO] A. Bondal, D. Orlov, Reconstruction of a variety from the derived category and groups of autoequivalences, alg-geom/97120029.
- [CDS] A. Connes, M. Douglas, A. Schwarz, Non-commutative geometry and Matrix theory: compactification on tori, hep-th/9711162.
- [Co] A. Connes, Non-commutative geometry.
- [CR] A. Connes, M. Rieffel, Yang-Mills for non-commutative two-tori. Contemporary Math. 62 (1987), 237-266.
- [De1] C. Deninger, On dynamical systems and their possible significance for arithmetic geometry. In: Regulators in analysis, geometry and number theory. Progress in mathematics 171, 29-87 (1999).
- [De2] C. Deninger, Number theory and dynamical systems on foliated spaces, preprint, 2001.
- [Di] R. Dijkgraaf, Mirror symmetry and elliptic curves. In: Moduli Space of Curves. Progress in Mathematics, vol. 129, p.149-163.

- [Ga] P. Gabriel, Des categories abeliennes. Bull. Soc. Math. France 90 (1962)
- [GLO] V. Golyshev, V. Lunts, D. Orlov, Mirror symmetry for abelian varieties, math.AG/9812003.
- [Ki] R. Kiehl, Theorem A und Theorem B in der nichtarchimedischen Funktionentheorie, Inv. Math. 2 (1967), 256-273.
- [Ko1] M. Kontsevich, Homological algebra of Mirror symmetry. Proceedings of the ICM in Zurich, vol. 1, p. 120-139.
- [Ko2] M. Kontsevich, Operads and Motives in Deformation Quantization, math.QA/9904055.
- [KR] M. Kontsevich, A. Rosenberg, Non-commutative smooth spaces, math.AG/9812158.
- [KoSo1] M. Kontsevich, Y. Soibelman, Homological mirror symmetry and torus fibrations, math.SG/0011041, Published in Proceedings of KIAS conference in Symplectic geometry and Strings, 2001.
- [KoSo2] M. Kontsevich, Y. Soibelman, Deformation theory (book in preparation).
- [M1] Yu. Manin, Quantized theta-function. Preprint RIMS, RIMS-700, 1990.
- [M2] Yu. Manin, Real multiplication and non-commutative geometry math.AG/0202109.
- [M3] Yu. Manin, Mirror symmetry and quantization of abelian varieties, math.AG/0005143.
- [M4] Yu. Manin, The theory of commutative formal groups over fields of finite characteristic. Russian Mathematical Surveys, Volume 18, n. 6.
- [Mo1] G. Moore, Finite in all directions, hep-th/9305139.
- [Mo2] G. Moore, Arithmetic and Attractors, hep-th/9807087
- [O1] D. Orlov, On equivalences of derived categories of coherent sheaves on abelian varieties, alg-geom/9712017
- [O2] D. Orlov, Equivalences of derived categories and K3 surfaces, alg-geom/9606006.

[PZ] A. Polishchuk, E. Zaslow, Categorical mirror symmetry: the elliptic curve, math.AG/9801119.

[RS] M. Rieffel, A. Schwarz, Morita equivalence of multidimensional non-commutative tori, math.QA/9803057.

[S1] A. Schwarz, Morita equivalence and duality, hep-th/9805034.

[S2] A. Schwarz, Theta-functions on noncommutative tori, QA/0107186.

[Sab] C. Sabbah, Systemes holonomes d'equations aux q-differences. In: D-modules and microlocal geometry, 1992, p. 125-147.

[So1] Y. Soibelman, Quantum tori, mirror symmetry and deformation theory, math.QA/0011162

Addresses:

Y.S.: Department of Mathematics, KSU, Manhattan, KS 66506, USA

email: soibel@math.ksu.edu

V.V.: Department of Mathematics, University of Chicago, Chicago, IL 60637

email: volgdsky@math.uchicago.edu